

nemaflor[®] - Biological control of Western Flower Thrips


nemaflor[®] contains 250 million infective juveniles of the insect pathogenic nematode *Steinernema feltiae* for biological control of *Frankliniella occidentalis* in greenhouses and tunnels.

Application conditions

For optimum results treat plants as soon as they are placed in the greenhouse.

nemaflor[®] does not control immigrant thrips. Most of these do not harm the crop.

Air temperature should be 10 to 28°C. At higher temperatures, nematode efficacy decreases. Nematodes are very sensible to sunlight and desiccation and should only be applied in the early morning or evening.




Application directions and rates

nemaflor[®] should be applied weekly as a routine application, since nematodes attack only the stages present at the time of application. The recommended application rate is 250,000 nematodes per m² per week or one pack of 250 million on 400 m² soil surface. Thrips numbers should be monitored weekly with blue or yellow sticky traps!

Foliar Application

Method recommended e.g. for Chrysanthemums. Apply 10-20 litres of water per 100 m². Spray to the point of run off. Apply in the evening to make use of the higher humidity. Growing points of plants should stay moist for at least 3 hours after application, to enable the nematodes to find their hosts.



Application via drip irrigation (Dosatron)

Method recommended for strawberries. Agitate suspension in the feeding tank continuously during injection to prevent nematodes from settling. At low infestation apply every two weeks with 250,000 nematodes per m².

Storage and handling

nemaflor[®] should be used as soon as possible upon delivery. Store at 2 to 8°C (see expiry date on package). It should never be frozen or exposed to more than 30°C. Remove all sieves in the tubing and at the nozzles. Use nozzles with at least 0.8 mm diameter. Do not exceed pump pressure of 5 bar. Mix the solution thoroughly and maintain agitation during spraying to avoid sedimentation of the nematodes!

Compatibility

IPM compatible insecticides have no effect on nematodes and can be used in rotation programs and as a tank mix partner. **nemaflor[®]** can be tank mixes with Naturalis (*Beauveria bassiana*).

Safety

The active ingredient and the formulation of **nemaflor[®]** are safe to users, consumers and the environment. There are no restrictions or safety instructions for the usage of **nemaflor[®]**.

