

nemastar[®]

For biological control of leatherjackets, mole crickets and cutworms

Storage

Apply nemastar[®] as soon as possible upon delivery. The product may be stored at 4-8°C until the expiration date on the package. Do not freeze or expose to temperatures higher than 35°C.

Application timing

Leatherjackets: Mid September to mid October
Mole Crickets: April to September
Cutworms: June to August

Application conditions

The soil temperature must be above 14°C for at least several hours per day. Apply during early morning or late evening. Do not apply in bright sunlight. The soil surface should be moist and remain moist for one week. Irrigate within 30 minutes after application to wash the nematodes into the soil.



Application/Dosage

Dissolve the package contents in a bucket of 10 litres of water and dilute to the desired amount of water. Rinse the package to flush any nematodes still clinging to it. Stir the solution until all lumps have dissolved. Repeat stirring every 10 minutes.

Package size	Premix for watering can in	Treats
250 million	50 litres nematode suspension	500 m ²
500 million	100 litres nematode suspension	1,000 m ²

Use the entire contents of the pack at once and completely use up the mixed spray liquid. Remove all filters and sieves and use nozzles with a diameter of at least 0.8 mm.

Stir during application to prevent the nematodes from settling.



Any questions?

beratung@e-nema.de

We can help you!

+49 4307/8295-295

www.e-nema.de