

nematop[®] Biological Soil Insecticide

For control of **vine weevil larvae and pupae** (*Otiorhynchus sulcatus*)

ACTIVE INGREDIENT: <i>Heterorhabditis bacteriophora</i>	15%
INERT INGREDIENTS.....	65%
WATER.....	20%

Bag holds 50 million *Heterorhabditis bacteriophora* nematodes. The nematodes are formulated in a wettable powder, enclosed in a Non-Water-Soluble pack.

**STORE AT 4° - 12° C
DO NOT FREEZE**

DIRECTIONS FOR USE

APPLICATION TIMING

Apply nematop from **end of March to the end of May** for treatment of larvae which have developed during winter and from **August to mid-October** against larvae which have developed from eggs laid in late spring.



CONDITIONS

- Apply nematop when soil temperatures are above 12° C for at least several hours a day.
- Apply early morning or late evening to minimize the effect of heat and sunlight.
- The soil surface should be moist at the time of application.
- Apply nematop with sufficient water (1-4 litres/m²) to moisten the soil surface.
- Irrigate within 30 minutes after treatment with at least 1 litre/ m² to wash nematop off the leaves and below the soil surface.
- For optimum results, maintain the soil moist for 3-4 weeks after application.

APPLICATION RATES

Soil: 0.5 million nematodes per 1 m².

Pots and Containers: 10,000 nematodes per litre potting compost.

Strawberry: 25,000 nematodes per plant.

- Use nematop as soon as possible upon delivery.
- Use the entire pack at one time.
- The entire spray solution should be applied once nematop is mixed with water. Do not store for longer than 4 hours.
- Remove all fine filters in the tubing and at the nozzles. Use nozzles with at least 0.8 mm diameter. Do not exceed a pump pressure of 5 bar.

APPLICATION

◆ Watering Can

- Dissolve one pack of nematop in 10 litres of water in a clean graduated bucket. Add one litre of the solution to a watering can and top up with water.



- 10 litres diluted solution in the watering can will treat 10 m².
- Repeat the above procedure 10x = 100 m² are treated.

◆ Ground Spraying Equipment

- Rinse equipment with water before using nematop.
- Add water to the spray or tank mix.
- With the agitator running, add the required amount of nematop through the sieve.
- Begin application of the spray solution and maintain agitation to avoid settling of the nematodes until all the mixture has been applied.



◆ Irrigation Equipment

- nematop can be applied through microirrigation (individual spaghetti tube), drip irrigation, or hand-held or motorised calibrated irrigation equipment.
- Irrigate prior to nematode application to moisten soil surface.
- Dissolve nematop in water and add it to a clean feeder tank. Agitate solution continuously during injection to prevent nematodes from settling. Alternatively add carboxy methyl cellulose (wallpaper paste) at 0.4% w/v.
- After injection, rinse feeder tank with clean water and inject into system.

◆ Dipping

- Add one pack to 80 litres of water and mix thoroughly the solution.
- Keep nematode solution agitated by bubbling in of compressed air.
- Alternatively add carboxy methyl cellulose (wallpaper paste) at 0.4% w/v.
- 80 litres will treat approx. 400 plants with fist- size root balls.

◆ Drench Application to Containers

Mix one pack of 50 million nematodes in 400 litres of water and evenly apply to surface of potting media. Adjust volume accordingly for smaller or bigger pots as outlined in the drench mixing chart.

Drench Mixing Chart

Pot Ø (cm)	Pot Volume (litres)	Drench per Pot (ml)	Pots per Pack
10	0.5	40	10,000
12	0.8	65	6,250
14	1.5	120	3,333
	3	240	1,667
	5	400	1,000

STRAWBERRY

◆ Dipping of Frigo Plants

- Add one pack to 8 litres of water and mix thoroughly the solution.
- Add carboxy methyl cellulose (wallpaper paste) at 0.4% w/v to prevent nematodes from settling.
- One frigo plant takes up approx. 2 ml of solution. 8 litres will treat 4,000 plants.

◆ Drench Application

- Drench 200 ml around the base of each plant or inject into the root zone.
- One pack of 50 million nematodes diluted in 400 litres of water will treat 2,000 plants.

COMPATIBILITY

nematop is compatible with most chemical fertilizers and pesticides. Only a few insecticides have a negative impact on nematop. In these cases apply nematop 3 days earlier or 2 weeks later. A compatibility list can be downloaded from www.e-nema.de.

STORAGE

The product may be stored for a limited period of time at 4 to 12°C (see expiring date on package).

SAFETY

The active ingredient and the formulation of nematop are safe to users, consumers and the environment. There are no restrictions or safety instructions for the use of nematop.